


List of Faunal species in and around greater Sipajhar Area

Teacher in-charge: Dr. Dibya Jyoti Saikia, HOD, Zoology Dptt.

- Collection of Photographs by:**
1. Kumar Kritartha Kaushik , Assistant Prof., Zoology Dept.
 2. Dhiraj Deka (Student)
 3. Pranjit Deka (Student)
 4. Bhargav Sandilya (Student)
 5. Subhrajit Saharia (Student)



4(Lemon pansy), 5(Chocolate pansy), 6(Leopard lacewing), 7(Leopard lacewing), 8(Blue spotted crow), 9(Grey count), 10(Common crow), 11(Great egg fly), 12(Common five ring), 13(Eastern five ring), 14(Common palm fly), 15(Common evening brown), 16(Common bush brown), 17(Common baron), 18(Knight), 19(Common sailor), 20(Large yeoman), 21(Small branded swift), 22(Coon), 23(Common banded demon), 24(Fulvous pied flat), 25(Grass demon), 26(Banded tree brown), 27(Peacock pansy), 28(Zebra blue), 29(Plain tiger), 30(Striped tiger), 31(Grey pansy)



Black-hooded oriole

Birds

Description

The black-hooded oriole is a member of the oriole family of passerine birds and is a resident breeder in tropical southern Asia from India and Sri Lanka east to Indonesia. It is a bird of open woodland and cultivation. The nest is built in a tree, and contains two eggs. [Wikipedia](#)

Scientific name: Oriolus xanthornus

Indian pied myna

Description

The Indian pied myna is a species of starling found in the Indian subcontinent. It is usually found in small groups mainly on the plains and low foothills. It is often seen within cities and villages although it is not as bold as the common myna. It produces a range of calls made up of liquid notes. [Wikipedia](#)

Scientific name: Gracupica contra



White-breasted waterhen

Birds

Description

The white-breasted waterhen is a waterbird of the rail and crane family, Rallidae, that is widely distributed across South and Southeast Asia. They are dark slaty birds with a clean white face, breast and belly. [Wikipedia](#)

Scientific name: Amaurornis phoenicurus





-Oriental magpie-robin

Birds

Description

The Oriental magpie-robin is a small passerine bird that was formerly classed as a member of the thrush family Turdidae, but now considered an Old World flycatcher. They are distinctive black and white birds with a long tail that is held upright as they forage on the ground or perch conspicuously. [Wikipedia](#)

Scientific name: Copsychus saularis

Red-throated barbet

Birds

Description

The red-throated barbet is a species of bird in the family Megalaimidae. It is found in Brunei, Indonesia, Malaysia, Myanmar, Singapore, and Thailand. Its natural habitats are subtropical or tropical moist lowland forest and subtropical or tropical swamps. It is threatened by habitat loss. [Wikipedia](#)

Scientific name: Megalaima mystacophanos



Black-rumped flameback

Description

The black-rumped flameback, also known as the lesser golden-backed woodpecker or lesser goldenback, is a woodpecker found widely distributed in the Indian subcontinent. It is one of the few woodpeckers that are seen in urban areas. It has a characteristic rattling-whinnying call and an undulating flight. [Wikipedia](#)

Scientific name: Dinopium benghalense



Italian sparrow

Birds

Description

The Italian sparrow, also known as the cisalpine sparrow, is a passerine bird of the sparrow family Passeridae, found in Italy and other parts of the Mediterranean region. [Wikipedia](#)

Scientific name: *Passer italiae*

Common myna

Birds

Description

The common myna or Indian myna, sometimes spelled mynah, is a bird in the family Sturnidae, native to Asia. An omnivorous open woodland bird with a strong territorial instinct, the common myna has adapted extremely well to urban environments. [Wikipedia](#)

Scientific name: *Acridotheres tristis*



Black drongo

Birds

Description

The black drongo is a small Asian passerine bird of the drongo family Dicruridae. It is a common resident breeder in much of tropical southern Asia from southwest Iran through India, Bangladesh and Sri Lanka east to southern China and Indonesia and accidental visitor of Japan. [Wikipedia](#)

Scientific name: *Dicrurus macrocercus*



Rufous treepie

Birds

Description

The rufous treepie is a treepie, native to the Indian Subcontinent and adjoining parts of Southeast Asia. It is a member of the crow family, Corvidae. It is long tailed and has loud musical calls making it very conspicuous. It is found commonly in open scrub, agricultural areas, forests as well as urban gardens. [Wikipedia](#)

Scientific name: *Dendrocitta vagabunda*



Rose-ringed parakeet

Birds

Description

The rose-ringed parakeet, also known as the ring-necked parakeet, is a medium-sized parrot in the genus *Psittacula*, of the family Psittacidae. [Wikipedia](#)

Scientific name: *Psittacula krameri*


Red-vented bulbul

Birds

Description

The red-vented bulbul is a member of the bulbul family of passerines. It is a resident breeder across the Indian subcontinent, including Sri Lanka extending east to Burma and parts of Bhutan and Nepal. [Wikipedia](#)

Scientific name: *Pycnonotus cafer*





Cattle egret

Birds

Description

The cattle egret is a cosmopolitan species of heron found in the tropics, subtropics, and warm-temperate zones. It is the only member of the monotypic genus *Bubulcus*, although some authorities regard its two subspecies as full species, the western cattle egret and the eastern cattle egret. [Wikipedia](#)

Scientific name: *Bubulcus ibis*


White-throated kingfisher

Birds

Description

The white-throated kingfisher also known as the white-breasted kingfisher is a tree kingfisher, widely distributed in Asia from the Sinai east through the Indian subcontinent to China and Indonesia. This kingfisher is a resident over much of its range, although some populations may make short distance movements. [Wikipedia](#)

Scientific name: *Halcyon smyrnensis*





Spotted dove

Birds

Description

The spotted dove is a small and somewhat long-tailed pigeon that is a common resident breeding bird across its native range on the Indian subcontinent and in Southeast Asia. The species has been introduced to many parts of the world and feral populations have become established. [Wikipedia](#)

Scientific name: *Spilopelia chinensis*

Lineated barbet

Birds

Description

The lineated barbet is an Asian barbet native to the Terai, the Brahmaputra basin to Southeast Asia. It is a frugivore and nests in holes of tree trunks. [Wikipedia](#)

Scientific name: *Megalaima lineata*



House crow

Birds

Description

The house crow, also known as the Indian, greynecked, Ceylon or Colombo crow, is a common bird of the crow family that is of Asian origin but now found in many parts of the world, where they arrived assisted by shipping. It is between the jackdaw and the carrion crow in size but is slimmer than either. [Wikipedia](#)

Scientific name: *Corvus splendens*